



Effect of Seaweed (*Gracilaria edulis*) Extract and Phosphorus on Growth and Economic of a Blackgram (*Vigna mungo* L.)

S. Sam Praveen Kumar^{a†}, Shikha Singh^{b#} and Neela Avani Pradeepika^{a†}

^a Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj-211007, Uttar Pradesh, India.

^b Naini Agricultural Institute, Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj-211007, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i1130933

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/85259>

Received 02 February 2022

Accepted 21 March 2022

Published 25 March 2022

Original Research Article

ABSTRACT

A field experiment trial on blackgram was conducted during *Zaid*, 2021 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P), India. The soil of the experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.3), organic carbon (0.73%), available nitrogen (274.46 kg/ha), available phosphorus (31.10 kg/ha), and available potassium (328 kg/ha). The foliar spray was applied thrice at different concentrations (5.0, 7.5, 10, and 15.0% v/v) of seaweed extract of *Gracilaria edulis* (*G. Sap*) with different levels of phosphorus (30 and 40 kg/ha) and control plot with water spray. The experiment was laid out in Randomized Block Design with nine treatments each replicated thrice. The results showed that *viz*: plant height (42.84 cm), number of branches per plant (7.07), number of nodules per plant (14.11), root dry weight (0.41 g/plant) were recorded significantly higher with 15% *G. Sap* + Phosphorus 40 kg ha⁻¹. Whereas, plant dry weight (17.55 g/plant) was recorded significantly higher with 10% *G. Sap* + Phosphorus 40 kg ha⁻¹. Also, the highest gross returns (62,716.50 INR/ha), net returns (26,259.00 INR/ha), and benefit-cost ratio (1.72) were obtained with the application of 15% *G. Sap* + Phosphorus 40 kg ha⁻¹ when compared to other treatments. Thus, foliar application of seaweed extract could be a promising option for yield enhancement in blackgram.

[†]Research Scholar;

[#]Assistant Professor;

*Corresponding author: E-mail: sampraveenkumar9@gmail.com;

Keywords: Seaweed extract; phosphorus; blackgram; growth and economic.

1. INTRODUCTION

Blackgram is scientifically known as *Vigna mungo* (L.) and commonly known as Urd in India. It is a tropical leguminous plant [1]. A family leguminoceae are includes about 18,000 species, which are characterized by their pods and alternate pinnate or trifoliate leaves Grain legumes, commonly known as pulse belong to sub-family papilionaceae as their flowers resemble the shape of butterfly [2]. It is popular because of its nutritional quality having rich protein (22-24%), carbohydrates (56.6-59.6%), fat (1.2-1.4%), Minerals (3.2%), phosphorous (385 mg/100g) and it is rich source of calcium and iron. It defers from other pulses in its peculiarity of attaining a somewhat mucilaginous pasty character, giving additional body to the mass due to long polymer chain of polysaccharide chain of carbohydrates. Tamil Nadu leads first in productivity with an average yield of 775 kg/ha. It contained, 24.7 % protein, 0.6 % fat, 0.9 % fibre and 3.7 % ash as well as sufficient quantity of calcium, phosphorus and important vitamins. Due to cheaper protein source of it is designated as "poor man's meat" [3].

Saline environmental bioactive algae extracted from seaweed are currently used in food, animal supplement, as raw materials in the industry. India is striving to increase its seaweed production to 11.2 lakh tonnes by 2025, up from the current production levels of 0.25 lakh tonnes in 2021 [4]. Marine bioactive substances extracted from seaweed play a great role in a new generation of natural organic fertilizer which contains all essential nutrients, amino acids, vitamins, cytokinin, and auxin-like promoting hormone. Seaweed act as a plant nutrient-bearing fertilizer. Especially in foliar spray, as a means to avoid excessive fertilizer application and to improve mineral absorption through leaves [5]. Using macroalgal agricultural bio-stimulants (ABs) on crop plants can generate numerous benefits with reported effects, including enhanced rooting, higher crop and fruit yields, enhanced photosynthetic activity, and resistance to fungi and bacteria viruses [6].

Application of phosphorus to pulse crop has been found very effective and called a master key element for increasing yield, it plays a vital role in the growth and development of roots and nodulation [7]. Judicious, use of phosphate fertilizer is supposed to result in better nodulation and efficient functioning of nodules bacteria for

fixation of atmospheric nitrogen to be utilized by the plant during the grain development stage, which in turn leads to an increase in grain yield [8]. It plays an important role in plants' main metabolic processes, including photosynthesis, energy transfer, signal transduction, macromolecular biosynthesis, and respiration.

2. MATERIALS AND METHODS

The method that was carried out for the germination test was "Top of Paper (TP) Media" at Agronomy laboratory, SHUATS, Prayagraj on 16th March 2021 [Fig.1]. Germination of blackgram *var. Shekhar-2* had recorded as 86.6%. A field trial was conducted during *Zaid*, 2021 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P), India which is located at 25°39'42" N latitude, 81°67'56" E longitude, and 98m altitude above the mean sea level (MSL). The soil was sandy loam in texture, low in organic carbon and medium in available nitrogen, phosphorus, and low in potassium. Nutrient sources were Urea, Single Super Phosphate, and Murate of potash to fulfill the requirement of Nitrogen, Phosphorus, and Potassium. The experiment was laid out in Randomized Block Design with nine treatments each replicated thrice [Fig.2]. The treatments which are with 1- 5% G. *Sap* + Phosphorus 30 kg ha⁻¹, 2- 5% G. *Sap* + Phosphorus 40 kg ha⁻¹, 3- 7.5% G. *Sap* + Phosphorus 30 kg ha⁻¹, 4 - 7.5% G. *Sap* + Phosphorus 40 kg ha⁻¹, 5 - 10% G. *Sap* + Phosphorus 30 kg ha⁻¹, 6 - 10% G. *Sap* + Phosphorus 40 kg ha⁻¹, 7- 15% G. *Sap* + Phosphorus 30 kg ha⁻¹, 8 - 15% G. *Sap* + Phosphorus 40 kg ha⁻¹, 9 - Control water spray (20:40:20 NPK kg/ha). Blanket application of a recommended dose of Nitrogen and Potassium (20:0:20 NPK kg/ha). The date of sowing was 24th march 2021 with the seed rate of 20kg/ha. "Line Sowing" method carried for sowing [Fig.3]. The foliar application seaweed (*Gracilaria edulis*) extract was sprayed at 15, 30 & 45 DAS [Fig.4]. The growth parameters of the plants were recorded at frequent intervals from germination up until harvest and finally, the yield parameters were recorded after harvest. The growth parameters reading such as plant height, number of branches, number of nodules, root dry weight, plant dry weight [Fig.5&6]. These parameters were recorded and statistically analyzed using analysis of variance (ANOVA) as applicable to Randomized Block Design [9]. Economic analysis is also calculated according to valid data.



Fig.1. Germination test of blackgram was done in the laboratory of the Department of Agronomy, SHUATS, Prayagraj



Fig. 2. The layout of the experimental field, Tilling the research plots through rake and leveling the plots through leveler at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj during Zaid, 2021



Fig. 3. The line sowing method was carried out manually for the sowing of blackgram seeds at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj during Zaid, 2021



Fig. 4. Weeding, thinning, and Foliar application of *G. Sap* (15DAS) was done manually at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj during *Zaid*, 2021



Fig. 5. Blackgram plant samples were uprooted at 40DAS for observation of dry matter accumulation and several nodules count



Fig. 6. Blackgram root dry weight was taken off accordingly and the Threshing of matured pods from the 1m² area had kept separately

3. RESULTS AND DISCUSSION

Effect on the growth and Growth attributes of blackgram. As can be seen in Table 1, growth parameters are summarized statistically. The Significantly taller plant height (42.84 cm) was recorded in treatment 8 with 15% *G. Sap* + Phosphorus 40 kg ha⁻¹. However, 10% *G. Sap* + Phosphorus 30 kg ha⁻¹ (40.60cm) and 10% *G. Sap* + Phosphorus 40 kg ha⁻¹ (41.24cm) statistically at par with 15% *G. Sap* + Phosphorus 40 kg ha⁻¹. The minimum plant height (35.85cm) was recorded in the treatment combination 7.5% *G. Sap* + Phosphorus 30 kg ha⁻¹. A significantly higher number of branches (7.07) recorded treatment 8 with 15% *G. Sap* + Phosphorus 40 kg ha⁻¹. However, 10% *G. Sap* + Phosphorus 40 kg ha⁻¹ (6.60) statistically at par with 15% *G. Sap* + Phosphorus 40 kg ha⁻¹. The minimum number of branches (5.87) were recorded in the treatment combination 5% *G. Sap* + Phosphorus 30 kg ha⁻¹. A maximum number of nodules (14.11) recorded treatment 8 with 15% *G. Sap* + Phosphorus 40 kg ha⁻¹. However, 5% *G. Sap* + Phosphorus 40 kg ha⁻¹ (12.78), 10% *G. Sap* + Phosphorus 40 kg ha⁻¹ (13.44) statistically at par with 15% *G. Sap* + Phosphorus 40 kg ha⁻¹. Significantly maximum root dry weight (0.41g) noted on 15% *G. Sap* + Phosphorus 40 kg ha⁻¹. However, 10% *G. Sap* + Phosphorus 40 kg ha⁻¹ (0.36) statistically at par with 15% *G. Sap* + Phosphorus 40 kg ha⁻¹. The minimum root dry weight (0.23 g/plant) was recorded with application of 5% *G. Sap* + Phosphorus 30 kg ha⁻¹ and 7.5% *G. Sap* + Phosphorus 30 kg ha⁻¹. The highest plant dry matter accumulated in treatment 6 with 10% *G. Sap* + Phosphorus 40 kg ha⁻¹ (17.55g). However, 7.5% *G. Sap* + Phosphorus 40 kg ha⁻¹ (17.47), 15% *G. Sap* + Phosphorus 30 kg ha⁻¹ (17.39) and 15% *G. Sap* + Phosphorus 40 kg ha⁻¹ (17.49) statistically at par with 15% *G. Sap* + Phosphorus 40 kg ha⁻¹. The minimum plant dry weight (17.16) was recorded with application of 5% *G. Sap* + Phosphorus 30 kg ha⁻¹. There was no significant difference in crop growth rate among the treatments at 40-60 DAS. maximum Crop growth rate (11.84 g/m²/day) were recorded in the treatment combination 7.5% *G. Sap* + Phosphorus 30 kg ha⁻¹ [Fig.7]. However, lowest (11.63 g/m²/day) was observed in 15 % *G. Sap* + Phosphorus 40 kg ha⁻¹. There was no significant difference in relative growth rate among the treatments at 40-60 DAS interval. The maximum relative growth rate were recorded in the treatment combination

of 5% *G. Sap* + Phosphorus 30 kg ha⁻¹ and 10% *G. Sap* + Phosphorus 30 kg ha⁻¹ which is 0.044 [Fig.8]. The minimum Relative growth rate were recorded in the treatment combination of 10% *G. Sap* + Phosphorus 30 kg ha⁻¹ and 10% *G. Sap* + Phosphorus 40 kg ha⁻¹ which is 0.041. The results demonstrate that the increase in plant height may be due to the seaweed saps which contain growth hormones like cytokinin and gibberellins that enhanced the growth of the plant. Similar results were reported that seaweed extract contains growth regulators like betaines and oligosaccharides that induced growth responses in the plants [10]. Seaweed extracts contain major and minor nutrients, amino acids, vitamins, cytokinins, auxin, and abscisic acid-like growth-promoting substances and have been reported to stimulate the height, branches, nodules, and yield of plants, develop tolerance to environmental stress [11]. This increase in growth parameters and yield might be due to phosphorus which is indispensable, a constituent of nucleic acid, ADP, and ATP. It has beneficial effects on nodulation, root development, growth, hastens maturity, and improves the quality of crop produce [12]. Extract of seaweed is known to contain nutrients and growth regulators such as auxins (IAA and IBA), gibberellins, cytokinins, betaines, and major macro and micronutrients. An increase in root length, root growth, number of lateral roots, and chlorophyll contents have been reported in different crops by seaweed extract application [13]. Maize dry matter accumulation significantly increased by increasing concentration of *Kappaphycus alvarezii* (*K. sap*) and *Gracilaria edulis* (*G. sap*) extract (Singh et al., 2015) [14]. The plants attained more vigor with phosphorus, due to adequate supply and availability of nitrogen, phosphorus, potassium, and spacing in balanced combination, resulting in increased dry weight of the plant. The application of Phosphorus to blackgram significantly increased dry matter production. [15] noted similar results. As phosphorus plays a critical role in various physiological processes, including root development, nodulation, and nitrogen fixation, better utilization of growth hormones, increased parameters may have been results of soil stimulation at 40kg/ha of phosphorus. Similar results were also observed by [16] & [17]. [18] experimented to study that the application of 40kg ha⁻¹ of phosphorus with spacing 30x10cm in blackgram had also shown almost similar results.

Table 1. Effect of Seaweed (*Gracilaria edulis*) Extract and Phosphorus on Growth of Blackgram

Treatment	At harvest				40-60 DAS		
	Plant height (cm)	Number of branches/ plant	Number of nodules/ plant	Root dry weight (g/plant)	Plant dry weight (g/plant)	CGR (g/m ² /day)	RGR (g/g/day)
1	38.05	5.87	10.77	0.23	17.04	11.80	0.044
2	39.27	6.07	12.78	0.29	17.13	11.66	0.042
3	36.36	6.27	9.78	0.23	17.18	11.84	0.044
4	37.61	6.27	11.44	0.32	17.47	11.73	0.042
5	40.60	6.13	10.78	0.26	17.19	11.64	0.042
6	41.24	6.60	13.44	0.36	17.55	11.63	0.041
7	38.55	6.13	10.77	0.31	17.39	11.75	0.042
8	42.84	7.07	14.11	0.41	17.49	11.63	0.041
9	38.05	6.00	11.66	0.24	15.92	11.66	0.042
F-test	S	S	S	S	S	NS	NS
SEm(±)	0.98	0.16	0.61	0.02	0.11	0.12	0.00
CD 5%	2.93	0.48	1.84	0.05	0.34	-	-

1- 5% G. Sap + Phosphorus 30 kg ha⁻¹, 2- 5% G. Sap + Phosphorus 40 kg ha⁻¹, 3- 7.5% G. Sap + Phosphorus 30 kg ha⁻¹, 4- 7.5% G. Sap + Phosphorus 40 kg ha⁻¹, 5- 10% G. Sap + Phosphorus 30 kg ha⁻¹, 6- 10% G. Sap + Phosphorus 40 kg ha⁻¹, 7- 15% G. Sap + Phosphorus 30 kg ha⁻¹, 8- 15% G. Sap + Phosphorus 40 kg ha⁻¹, 9 - Control water spray (20:40:20 NPK kg/ha)




Fig. 7. Crop Growth Rate




Fig. 8. Relative Growth Rate

Table 2. Effect of Seaweed (*Gracilaria edulis*) Extract and Phosphorus on Economics of Blackgram

Treatment	Cost of cultivation (INR/ha)	Gross returns (INR/ha)	Net returns (INR/ha)	B: C ratio
1	32,832.00	46,193.49	13,361.49	1.41
2	33,457.00	50,847.93	17,390.93	1.52
3	33,582.00	49,108.50	15,526.50	1.46
4	34,207.00	54,700.38	20,493.38	1.60
5	34,332.00	47,338.83	13,006.83	1.38
6	34,957.00	56,972.79	22,015.79	1.63
7	35832.00	50,234.94	14,402.94	1.40
8	36,457.00	62,716.50	26,259.50	1.72
9	31,957.00	45,568.53	13,611.53	1.43

1- 5% *G. Sap* + Phosphorus 30 kg ha⁻¹, 2- 5% *G. Sap* + Phosphorus 40 kg ha⁻¹, 3- 7.5% *G. Sap* + Phosphorus 30 kg ha⁻¹, 4- 7.5% *G. Sap* + Phosphorus 40 kg ha⁻¹, 5- 10% *G. Sap* + Phosphorus 30 kg ha⁻¹, 6- 10% *G. Sap* + Phosphorus 40 kg ha⁻¹, 7- 15% *G. Sap* + Phosphorus 30 kg ha⁻¹, 8- 15% *G. Sap* + Phosphorus 40 kg ha⁻¹, 9 - Control water spray (20:40:20 NPK kg/ha)
 *Economics not subjected to data analysis

Economic analysis of blackgram. As can be seen in Table 2, all the economic parameters viz. cost of cultivation, gross return, the net return, and B: C ratio are summarized. Cost of cultivation (36,457.00 INR/ha) was found to be highest in 15% *G.Sap* + Phosphorus 40 kg ha⁻¹ and the minimum cost of cultivation (31,957.00 INR/ha) was found to be in treatment with Control plot with water spray (20:40:20 NPK kg/ha) as compared to other treatments. The maximum gross return was observed in the treatment combination of blackgram was 15% *G.Sap* + Phosphorus 40 kg ha⁻¹ (62,716.50 INR/ha). The maximum net return was observed in the treatment combination of blackgram was 15% *G.Sap* + Phosphorus 40 kg ha⁻¹ (26,259.50 INR/ha). The maximum benefit cost ratio was observed in the treatment combination of blackgram was 15% *G.Sap* + Phosphorus 40 kg ha⁻¹ (1.72). The higher economic parameter was recorded with higher concentration and doses of treatment. Application of seaweed extracts and

RDF increased the gross and net returns as compared to sole RDF application of control plot. These results are in accordance with earlier findings of (Narendra et al., 2018) [19] and (Nayak et al., 2020) [20].

4. CONCLUSION

Based on my research trail, the treatment combination of 15% *G. Sap* with phosphorus 40kg/ha was found to be more productive and also economically feasible. Although the findings are based on one season, further research is needed to confirm the findings and their recommendation.

ACKNOWLEDGEMENT

The authors are thankful to Department of Agronomy, Naini Agricultural Institute (NAI), Sam Higginbottom University of Agriculture,

Technology and Sciences, Prayagraj- 211007, Uttar Pradesh, India for providing field, necessary facilities and assistance in conducting this research. Also, I would like to thank to the field workers of my department.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Rajeshkumar S, Nalliah Durairaj S, and Kannan V. Effect of Crop Geometry and Foliar Nutrition on Growth and Yield of Irrigated Blackgram, International Journal of current microbiology and applied sciences, 2017;6(11): 4084 -4094.
- Arulbalachandran D, Shankar Ganesh K, Subramani A. American Eurasian Journal of Agronomy. 2009;2(2):109-116.
- Aslam M, Hussain N, Zubair M, Hussain SB, and Baloch MS. Integration of organic and inorganic sources of phosphorus for increased productivity of mung bean (*Vigna radiata* L). Pakistan Journal of Agriculture science. 2010;47(2):111-114.
- FAO. 2020. The state of world fisheries and aquaculture (SOFIA-2020), Rome, Italy.
- Sam Praveen Kumar S, Avani Pradeepika N, and Divyalatha R. Seaweed: Farming and Uses of Extract in Agriculture. Agriculture & Food: E-Newsletter. 2022;4(2):74-75.
- Sharma HSS, Fleming C, Selby C, Rao JR, and Martin T. Plant biostimulants: a review on the processing of macroalgae and use of extracts for crop management to reduce abiotic and biotic stresses. Journal of Applied Phycology. 2014;26:465-490.
- Zhang X, Ervin EH, and Schmidt ER. Plant growth regulators can enhance the recovery of Kentucky bluegrass sod from heat injury. Crop Science. 2003;43:952-956.
- Dharwe DS, Dixit HC, Dotaniya CK, Douthaniya RK, Sandeep M, and Tarwariya, MK. Effect of Phosphorus and Sulphur on Yield Attributes, Nutrient Content and Nutrient Uptake of Green Gram in Bundelkhand Soil. International Journal of Current Research. 2019;11(11):8225-8229.
- Gomez, K.A., and A.A. Gomez. Statistical procedures for agricultural research. John Wiley and Sons, New York, 1984;680p.
- Balakrishnan CP, Kumar V, Mohan VR, and Athiperumalsami T. Study on the effect of crude seaweed extracts on seedling growth and biochemical parameters in *Cyamopsis tetragonoloba* (L.) Taub. Plant Archives. 2007;7(2):563-567.
- Yadav M, Yadav SS, Kumar S, Yadav T, and Yadav HK. Effect of Phosphorus and Bio-fertilizers on Growth and Yield of Urdbean [*Vigna mungo* L.]. International Journal of Plant & Soil Science. 2017;18(5):1-7.
- Choudhary R, Singh K, Manohar RS, Yadav AK, and Sangwan A. Response of different sources and levels of phosphorus on yield, nutrient uptake, and net returns on mung bean under rainfed conditions. Indian Journal of Agriculture Research. 2015; 35(4):263-268.
- Jadhao GR, Chaudhary DR, Khadse VA, and Zodape ST. Utilization of seaweeds in enhancing productivity and quality of blackgram (*Vigna mungo* L. Hepper) for sustainable agriculture. Indian Journal of Natural Products and Resources. 2015;6(1):16-22.
- Singh Shikha, Singh MK, Singh A, Singh CS. Application of seaweed sap (*Kappaphycus alvarezii* and *Gracilaria edulis*) for higher productivity of Maize (*Zea mays* L.) Research journal of Agricultural Sciences. 2015; 6(1):232-234.
- Kumar S, Tomar S, and Tomar TS. Integrated phosphorus management in blackgram (*Vigna mungo*) in western Uttar Pradesh during the summer season. Annals of Agricultural Research. 2014;35(3):290-297.
- Kadam SS, and Khanvilkar SA. Effect of phosphorus, boron, and row spacing on the growth of summer green gram (*Vigna radiata*). Journal of Agriculture and Crop Science. 2015;2:7-8.
- Sathyamoothi K, Mohamed, Amanullah E, Pazhanivelan S, and Vaiyapun K. Root growth and yield of a Greengram [*Vigna radiata* L.] as influenced by increased plant density and nutrient management. Journal of Applied Sciences Research. 2008;1(7): 917-924.
- Bonepally R, Umesha C, and Meshram MR. Influence of Spacing and Phosphorus Levels on Growth, Yield and Economics of

- a Blackgram (*Vigna mungo L.*). Biological Forum – An International Journal. 2021;13(1):82-85.
19. Narendra K, Mahendra S, Tomar IS, Dudwe TS, Yadav RK, and Yogesh KS. Effect of gibberellic acid on growth and yield and economic of Maize (*Zea mays L.*) Journal of Agricultural Research 2018;5(1):25-29.
20. Nayak P, Biswas S, and Dutta D. Effect of Seaweed extracts on growth, yield and economics of kharif rice (*Oryza sativa L.*). Journal of Pharmacognosy and Phytochemistry. 2020;9(3):247-253.

© 2022 Sam et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/85259>