



Economic Study of Resource Management Silver Pomfret (*Pampus argentus*) and Barramundi (*Lates calcarifer*) in Cirebon District, West Java - Indonesia

Dayang Diah Fidhiani ^{a*} and Dori Rachmawani ^b

^a Program Study of Aquaculture, School of Fisheries Science Malang, East Java, Indonesia.

^b Program Study of Aquatic Resources, Faculty of Fisheries and Marine Sciences University of Borneo Tarakan, North Kalimantan, Indonesia.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJFAR/2023/v21i6556

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97888>

Original Research Article

Received: 18/01/2023

Accepted: 23/03/2023

Published: 28/03/2023

ABSTRACT

The result catches Fish of Silver pomfret (*Pampus argentus*), and Barramundi (*Lates calcarifer*) contributed significantly to improving the welfare of Fishermen, especially Cirebon Regency of West Java. The study aimed to estimate the optimal utilization and formulate Fisheries Management Models. To pass the estimation and management model, used the Model Bioeconomy. The results obtained from the estimated balance of the Bioeconomy optimum utilization of fishery resources fish of Barramundi the Harvest of 707,18 kg/year, efforts of 7063,49 trips/year, and Phi of Rp.15.824.173, 45/year and fish of Silver pomfret the Harvest of 280,96 kg/year, efforts of 2660,20 trips/year, and phi of Rp.2.400.882,73/year and then be created for the

*Corresponding author: Email: fidhiani30@gmail.com;

formulation of Economic Models of Resource Management fish of Silver pomfret and Barramundi optimal and sustainable in Cirebon Regency is the management regime based on sole owner/Maximum Economic Yield (MEY) to form the optimal management of effort and catch.

Keywords: Management economical; fish; Cirebon regency.

1. INTRODUCTION

One of the potential marine resources that have been long exploited by the population is the fisheries resources [1]. Gross regional domestic product (GRDP) of the fishery sector in 2013-2019 has experienced an increasing growth every year. The calculation results show the value of the growth index of 177.22. It can be said the GRDP of the fisheries sector in Cirebon Regency has increased by 77.22% in the five year period [2]. Silver pomfret and Barramundi are one of the Demersal Fish groups that are mostly caught by Fishermen in Cirebon Regency and have economic value. According to Genisa (1999) in Mayalibit et al. [3], economically significant means having a high market value, tall and wide macro production volume, and increased production power. This activity allows for high profits, considerable resource potential, suitable water areas for these types of fish, and also due to the excellent market share of these fish. This opportunity has boomeranged for sustainable capture fisheries activities. Increasing fishing activity through increasing fishing efforts in terms of increasing the number of units and fishing trips will put great pressure on resources and ecosystems, leading to biologically and economically overfishing (biological and economic overfishing) [4].

Biological overfishing will reduce the amount of biomass (stock) which is characterized by a decrease in the number of catches per unit, changes in the size of the fish caught that are getting smaller, and changes in the fishing grounds (fishing ground). This biological decline will cause a decrease in economic terms, with reduced yields of catches on the one hand and an increase in the Cost of catches per trip; on the other hand, then fishing activities will lead to a breakeven point, where the costs incurred are equal to the results achieved and eventually will experience losses. This condition becomes inefficient, so an accurate estimate is needed [5].

This study aims to: (1) Determine the optimal utilization of White Pomfret and White Snapper fisheries in the Cirebon Regency, and (2) Formulate an optimal and sustainable resource management model.

2. RESEARCH METHODS

The research was conducted in Cirebon District, West Java Province. The data to be used in this study are primary and secondary data, including actual production, sustainable production, effort, CPUE, fishing costs, and fishing fleets.

2.1 Catch per Unit Effort

Catch per unit effort (CPUE) is calculated based on the total Catch of one fishing gear with the total effort of Catch (effort) in units. CPUE is calculated by the following formula [6] namely:

$$CPUE_{it} = Catch_{it} / Effort_{it}$$

Information

$CPUE_{it}$ = Catch per unit Effort of the time-t catching device

$Catch_{it}$ = The Catch (Catch) of fishing gear-i at time-t

$Effort_{it}$ = catching effort (effort) of fishing gear-i at time-t

2.2 Effort Standardization

Considering the diversity of fishing gear operating in the research area, to measure with equal units, effort is standardized between gear using the following standardization techniques developed by King [7], namely:

$$E_{it} = \varphi_{it} D_{it}$$

$$\text{With: } \varphi_{it} = U_{it} : U_{std}$$

Information

E_{it} = effort of standardized fishing gear

D_{it} = number of fishing days for fishing gear-i at time-t

φ_{it} = fishing power value of fishing gear-i in period-t

U_{it} = Catch per unit effort (CPUE) of fishing gear-i in period t

U_{std} = Catch per unit effort (CPUE) of the fishing gear used as the basis for standardization

2.3 Cost Standardization

Standardized Cost per unit effort (standardized unit effort) follows the standardization pattern used by Anna [8], which can be written mathematically as follows:

$$C_{et} = \left[\frac{1}{n} \sum_{i=1}^n \frac{TC}{\sum E_i} \left(\prod_{i=1}^n \frac{h_{it}}{\sum (h_i + h_j)} \right)^{\frac{1}{i-1}} \frac{CPI_t}{100} \right]$$

Information

C_{et} = Cost per unit of standardized effort in period t
 T_{ci} = total Cost for fishing gear i for $i = 1, 2$
 E_i = total standardized effort for gear i
 h_{it} = production of gear i at time t
 $\sum (h_i + h_j)$ = total fish production for all fishing gear
 n = number of fishing gear
 CP_{it} = consumer price index in period t

2.4 Estimation of Biological Parameters

Estimating the parameters r , q , and K for the two previous equations uses a non-linear technique, so by using the weighted least squares (WLS) method, namely by dividing the function $h(q, K, E)$ by $E(U_t = h_t / E_t)$, then these two equations can be transformed into linear equations so that the ordinary least squares (OLS) regression method can be used to estimate the biological parameters of the function mentioned above, namely as follows [8]:

Bentuk Gompertz : $\frac{h_t}{E_t} = qKe^{\left[\frac{-qE}{r}\right]}$

In $U_t = \beta_0 - \beta_1 E_t$, dimana $U_t = \frac{h_t}{E_t}$; $\beta_0 = \ln qK$; $\beta_1 = \frac{q}{r}$

The biological parameter estimation approach for sustainable production using the exponential-based Gompertz function is carried out using the parameter estimation model developed by Clarke, Yoshimoto, and Pooley [9] in Fauzi [5], namely as follows:

$$\frac{U_{t+1}}{U_t} - 1 = r - \frac{r}{qK} U_t - qE_t$$

$$\ln(U_{t+1}) = \frac{2r}{(2+r)} \ln(qK) + \frac{(2-r)}{(2+r)} \ln(U_t) - \frac{q}{(2+r)} (E_t + E_{t+1})$$

From the regression results, the value of β_1 will be obtained, which will justify the value of the

parameter r . In addition, the value of β_2 is also brought, which will justify the parameter value q . Furthermore, the K parameter values can be obtained from these values (r and q). The following equation is a form of the Clarke, Yoshimoto, and Pooley (CYP) [9] model approach to estimate biological parameters, namely:

$$r = \left(\frac{2 - 2\beta_1}{1 + \beta_1} \right)$$

$$q = \beta_2 (2 - r)$$

$$K = \frac{e^{\frac{\beta_0(2+r)}{2r}}}{q}$$

Information

K carrying capacity of the environment

q = capture coefficient
 r = intrinsic growth
 β_1 = intercept
 β_2 = slope

2.5 Estimation of Economic Parameters

Economic parameter estimation consists of price per kilogram, Cost per fishing trip, and discount rate. Prices and costs are measured in absolute terms adjusted for the consumer price index (CPI), where these values are obtained from nominal prices in period t (p_{nt}) and converted to accurate prices (p_{rt}). In more detail in the following equation:

$$p_{rt} = \left(\frac{p_{nt}}{IHK_t} \right) \times 100$$

Information

P_{rt} = accurate price in the year (t)
 P_{nt} = nominal price in the year (t)
 CP_t = consumer price index in the year (t)

While the discount rate is the real interest rate which is calculated based on the market base discount rate, and the interest rate calculation uses the following equation [10]:

$$\delta = \ln(1+r)$$

Information

δ = real discount rate
 r = interest rate

2.6 Bioeconomic Balance Estimation

The value of resource optimization using a static approach for bioeconomic equilibrium conditions in the maximum sustainable yield regime is obtained through the following equations [5], namely:

$$x_{MSY} = K \left(1 - \frac{qE}{r} \right)$$

$$E_{MSY} = \frac{r \left(1 - \frac{x}{K} \right)}{q}$$

$$h_{MSY} = qKE \left(1 - \frac{qE}{r} \right)$$

Information

x_{MSY} = biomass level
 h_{MSY} = optimal production rate
 E_{MSY} = optimal effort level
 K = carrying capacity of the environment
 q = capture coefficient
 r = intrinsic growth
 E = level of effort

Bioeconomic balance in conditions of open access can be denoted as follows:

$$x_{OA} = \frac{c}{pq}$$

$$h_{OA} = \frac{rc}{pq} \left[1 - \frac{c}{pqK} \right]$$

$$E_{OA} = \frac{r}{q} \left[1 - \frac{c}{pqK} \right]$$

Information:

x_{OA} = biomass level
 h_{OA} = optimal production level
 E_{OA} = optimal effort level
 K = carrying capacity of the environment
 q = capture coefficient
 r = intrinsic growth
 p = price of fish
 c = Cost of catching

For the sole owner condition, optimization is obtained based on the following equations:

$$x_{SO} = \frac{K}{2} \left(1 - \frac{c}{pqK} \right)$$

$$h_{SO} = \frac{rK}{4} \left[1 + \frac{c}{pqK} \right] \left[1 - \frac{c}{pqK} \right]$$

$$E_{SO} = \frac{r}{2q} \left[1 - \frac{c}{pqK} \right]$$

Information:

x_{SO} = biomass level
 h_{SO} = optimal production level
 E_{SO} = optimal effort level
 K = carrying capacity of the environment
 q = capture coefficient
 r = intrinsic growth
 p = price of fish
 c = Cost of catching

Furthermore, by using a dynamic approach, the optimization of fishery resources can be obtained with the following formula [5], namely:

$$x^* = \frac{1}{c} x (pqx - c) \left(\delta - r \left(1 - \frac{2x}{K} \right) \right)$$

$$h^* = \frac{1}{4} \left[\left(x_{OA} + K \left(1 - \frac{\delta}{r} \right) \right) + \sqrt{\left(x_{OA} + K \left(1 - \frac{\delta}{r} \right) \right)^2 + \left(\frac{8Kx_{OA}\delta}{r} \right)} \right]$$

$$E^* = \frac{h^*}{qx^*}$$

Information:

x^* = biomass level
 h^* = optimal production rate
 E^* = optimal effort level
 K = carrying capacity of the environment
 x_{OA} = biomass in open access conditions
 p = price of fish
 c = Cost of catching
 q = capture coefficient
 r = intrinsic growth
 δ = real discount rate

3. RESULTS AND DISCUSSION

3.1 Catch per Unit Effort

Catch per unit effort is a basic fisheries science often used as a first-order evaluation of board trends in likely relative abundance or relative

biomass trends over time of the underlying fish stocks [11-13]. While CPUE has its limitations, it can be useful in long term trend monitoring of a fishery, particularly where more detailed data and stock assessments do not exist (Stamatopoulos, 2022) and open or semi open access fisheries predominate, i.e., where restrictive fisheries management actions may be absent or ineffective [13]. Based on the catch per unit effort analysis results for Silver pomfret, it was found the highest CPUE value was the payang gear. This value illustrates the effectiveness of catching payang fishing gear, which is quite reasonable compared to other fishing gear in white pomfret fishing activities in Cirebon Regency waters. The opposite condition occurs for gill net fishing gear in white pomfret fishing activities, with an average CPUE value of 0.001560, indicating low fishing effectiveness. The results of the standardized effort analysis show that the standard fishing gear for catching the Silver pomfret in the waters of Cirebon Regency is the payang gear. Based on the effort of the standard fishing gear, the total effort for catching white pomfret is obtained. The following shows the actual effort and production level of Silver pomfret.

Furthermore, analysis of the Catch per unit effort of Barramundi found that longline fishing gear is the fishing gear that has the highest CPUE value, which is an average of 0.009834 per year. This value describes the ratio between Catch and effort. The higher the value, the better the effectiveness of the fishing gear. In other words, with fewer fishing efforts, more catches are obtained. The average longline effort is 10,167.27 trips per year, while the average gill net effort is 332,559.07.

Based on Fig. 1 it can be seen that between CPUE and fishing effort for Silver Pomfret in Cirebon Regency shows a negative relationship, meaning that the higher the fishing effort, the lower the CPUE value. The negative relationship indicates that the productivity of fishing gear decreases with increasing fishing effort, the lower the CPUE value. The negative relationship indicates that the productivity of fishing gear decreases with increasing fishing effort, the rate of resource exploitation leads to overfishing if not controlled [14,15]. Stock biomass is a limited resource jointly pursued by vessels in a fishery where the smaller the share per vessel, the more vessels that enter the fishery. Furthermore, the relationship between CPUE and fishing effort for Barramundi in Cirebon District shows a positive relationship, meaning the higher the fishing effort, the higher the CPUE value. The positive relationship indicates that the productivity of fishing gear increases with increasing fishing effort. Increasing and decreasing CPUE value also have a linear relationship with the density and availability of fish stocks in a water [14].

3.2 Bioeconomic Balance Estimation

Bioeconomics is a field of science that integrates biological and ecological resources with the economic behavior of fisherman, within consideration the dimension of space, time, and uncertainty. The relative importance of including some or all of the dimensions mentioned in the fisheries bioeconomic modelling and analysis will depend on the question related to fisheries management, stock mobility, and sensitivity to environmental factors and possible behavior of fisherman [16,17].




Fig. 1. Graph of the relationship between Catch Per Unit Effort (CPUE) and trips for silver pomfret and barramundi

The result of standardization of effort is that the payang is the standard fishing gear in white pomfret fishing activities in the waters of Cirebon Regency. The average effort is 16,100 trips per year, with an average catch of 937.50 tonnes per year. An average CPUE value of around 0.031667 per year is obtained with the average annual Catch an average effort. Furthermore, the regression analysis results obtained the value of $\beta_1 = 0.7492$ and $\beta_2 = 6.19101E-05$ with a value of $R^2 = 0.6391$. Knowing the value of the bioeconomic balance, the results of the analysis of Biological parameters and Economic parameters are simulated, and the optimal Catch (Harvest), optimal effort (Catch), and optimal rent (ϕ) values will be obtained. Based on the results of the Gordon-Schaefer (GS) Bioeconomic balance analysis, three bioeconomic balance points were obtained, namely the balance that occurs at the effort level of 6,709 trips per year, the effort level of 7,063 trips per year and an effort level of 13,419 trips per year. The first bioeconomic balance occurs in managing the Maximum Economic Yield (MEY) regime, commonly known as the sole owner. MEY's bioeconomic balance occurs at an effort level of 6,709 trips per year, with an optimal production level from an economic standpoint of around 17,593.69 tons per year. This condition will provide the highest profit in the form of resource rent of IDR 318,324,363 per year.

The second bioeconomic balance occurs in Maximum Sustainable Yield (MSY) resource exploitation conditions, commonly known as sustainable potential. The bioeconomic balance of the MSY regime occurs at an effort level of 7,063 trips per year, with an optimal catch of 17,637.89 tons per year. This condition will provide benefits in the form of resource rents of around IDR 317,440,384 per year. Thus if the resource utilization activities

Silver Pomfret in Cirebon Regency want to be sustainable from an ecological point of view, so input control must be carried out at a level of 7,063, which means reducing the current input of around 7,936.51 trips per year. These inputs can only be controlled by exploiting resources in the MEY and MSY regimes. These two regimes are forms of optimal management of resources from an economic, ecological, and social perspective (economic ecology and social sustainability). The third bioeconomic equilibrium occurs when the total cost (TC) curve intersects with the total revenue (TR) curve. This point is called the breakeven point, where $TR = TC$. Thus, under these conditions, no profit or resource rent equals zero ($\pi = 0$). Gordon calls this balance point the bioeconomic equilibrium of open access or the bioeconomic balance of open access [18]. This happens because, in open access conditions with no rules in utilization, each increase in effort will increase the total Cost of production so that the Total Cost is greater than the Total Revenue. Thus the exploitation of Silver Pomfret Fish Resources will suffer losses so that there will be input exits from resource actors, and the level of effort will return to the breakeven point.

Concerning the breakeven point of the bioeconomy balance, the exploitation of Silver Pomfret resources in the Cirebon Regency is currently experiencing overfishing and inefficiency. This can be seen in the bioeconomic balance Fig. 2a, with an actual effort value of 16,100 trips per year, exceeding the effort level at the breakeven point of 14,126 trips per year. This situation illustrates that Silver Pomfret fishery business activities will experience a reduction in inputs due to the large production costs incurred.




Fig. 2. Bioeconomic balance (a) Silver Pomfret (*Pampus argenteus*) and (b) Barramundi (*Lates calcarifer*)

Furthermore, the analysis of the bioeconomic balance in Fig. 2b shows that the management of Barramundi in the waters of Cirebon Regency from the point of view of Catch (Harvest) is experiencing overfishing. This can be seen from the research results, which show that the current catch value (68.70 tons per year) is far below the optimal catch value (6,269.38 tons per year). This value is unbalanced when looking at the number of actual efforts (24,084 trips), far from the optimal effort level (2,485). This condition illustrates that fishing activities for Barramundi in Cirebon Regency waters are inefficient. A very high amount of effort does not give optimal catch results. If this condition is maintained, it is believed that the Barramundi fishery business activity in the waters of Cirebon Regency will not last long. This can happen because of the amount of pressure due to the amount of effort that is too large. While the optimal effort value can be recommended so that Barramundi fishing activities can be sustainable from an ecological perspective, namely 2,660 trips per year, and from an economic perspective, namely 2,485 trips per year.

To achieve sustainable utilization of Barramundi resources in Cirebon Regency, it is necessary to reduce the amount of input (effort) by around 21,424 trips per year. So that the negative excesses of reducing the amount of effort can be maintained with the assumption that each operation of longline fishing gear requires a minimum of 2 crew members (referring to the results of the survey, 2007), there will be a reduction in the number of fishery actors by around 42,848 people. Alternative efforts to manage resources must always be carried out (back-forward looking), such as changing fishing gear with different catch targets with the requirement that the fishing gear targets are not optimal or diverting activities in other forms either in the fisheries sector such as marine and coastal cultivation or product processing, as well as in supporting fisheries activities based in coastal areas. This is in line with Adrianto [19] that the core of the transfer (transform) aims to move fishermen's livelihoods vertically and horizontally.

However, the transfer of profession must still pay attention to the characteristics and culture of the community (fishery actors). According to Charles (2001) in Adrianto [19] that there are several general characteristics of fishermen (fishermen), namely that first, fishermen differ according to social backgrounds such as age level, education,

social status, and level of cohesion in micro-communities (between fishermen in one group) or the macro community (fishermen with other coastal community members). Second, in the commercial fishing community, fishermen can vary according to their occupational commitment, such as full-time fishermen, main part-time fishermen, and part-time fishermen, or according to their occupational pluralism, such as fishermen with particular specialties, fishermen with various sources of income, and so on. Third, fishermen can vary according to motivation and behavior, where maximizers are fishermen who actively catch fish to get the maximum profit and tend to behave like a "company" and groups of fishermen satisficers or fishermen who actively catch fish to get enough results [20-28].

4. CONCLUSIONS AND RECOMMENDATIONS

4.1 Conclusion

1. Obtained optimal utilization of Barramundi (*Lates calcarifer*) resources, namely Harvest of 707.18 kg/year, effort of 7,063.49 trips/year, and Phi of Rp. 15,824,173.45/year and Silver Pomfret (*Pampus argentus*), namely Harvest of 280.96 kg/year, effort of 2,660.20 trips/year, and phi of IDR 2,400,882.73/year.
2. The formulation of an optimal and sustainable economic model for managing pomfret and Barramundi resources in Cirebon Regency is management based on the sole owner/Maximum Economic Yield regime, with an optimal form of effort and catch management.

4.2 Recommendations

Suggested that the regional government can regulate fishing efforts from exploiting barramundi fish to optimal exploitation levels, at MSY/MEY utilization levels so that fishery resources can be sustainable

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Yusrizal Wiyono ES, Simbolon D, Solihin I. Estimation of the Utilization Rate of Fish

- Resources in the Northern Coast of Java, Indonesia. *AAFL Bioflux*. 2018;11(6):1807-1824.
2. Harsen Faris, Rizal Achmad, Syamsudin Mega, Suryana Asep. The Contribution of Isheries Sector In Regional Development of Cirebon Regency of West Java Province, Indonesia. *Asian Journal of Fisheries and Aquatic Research*. 2021:1-12.
DOI: 10.9734/ajfar/2021/v13i130243
 3. Mayalibit DNK, Rahmat Kurnia, Yonvitner. Bioeconomic Analysis for Resource Management of Yellow Trevally (*Selaroides Leptolepis*, Cuvier & Valenciennes) Landed at PPN Karangantu, Banten. *Bonorowo Wetlands*. 2014;4(1):49-57.
 4. Fauzi A. *Ekonomi Perikanan: Teori, Kebijakan dan Pengelolaan*. Gramedia Pustaka Utama, Jakarta. 2010:245.
 5. Fauzi A. *Economics of Natural Resources and the Environment*. PT. Main Library Gramedia. Jakarta. 2004:259.
 6. Sparre Pdan, Venema SC. *Introduksi Pengkajian Stok Ikan Tropis*. Pusat Penelitian dan Pengembangan Perikanan: Jakarta; 1999.
 7. King M. *Fisheries biology, Assessment and NManagement*. London (UK): Fishing News Books; 1995.
 8. Anna S. *Model Embadded Dinamik Ekonomi Interaksi Perikanan-Pencemaran: Kasus di Teluk Jakarta DKI Jakarta*. Disertasi. Program Pasca Sarjana Institut Pertanian Bogor. 2023:371.
 9. Clarke RP, Yoshimoto SS, Pooley SG. A Bioeconomic Analysis of the North-Western Hawaiian Island Lobster Fishery. *Marine Resource Economics*. 1992;7(2):115-140.
 10. Kula E. Derivation of Social Time Preference Rates for the U.S and Canada. *Quarterly Journal of Economics*. 1984;99:873-882.
 11. Skalski JR, Ryding KE, Millspaugh JJ. *Wildlife Demography: Analysis of Sex, Age, and Count Data*. Elsevier Academic Press, Burlington, USA; 2005.
 12. Belhabib D, Lam VWY, Cheung WWL. Overview of West African fsheries under Climate Change: Impacts, Vulnerabilities and Adaptive Responses of the Artisanal and Industrial Sectors. *Mar. Policy*, 2016;71:15-28.
 13. Zeller D, Vianna GMS, Ansell M, Coulter A, Derrick B, Greer K, Noël S-L, Palomares MLD, Zhu A, Pauly D. Fishing Effort and Associated Catch per Unit Effort for Small-Scale Fisheries in the Mozambique Channel region: 1950–2016. *Front. Mar. Sci*. 2021;8:707999.
DOI: 10.3389/fmars.2021.707999
 14. Suman Rani, Raj Kumar Duhan, Sandeep Deshwal. Study of impementation of Total Quality Management for Defect Reduction in Manufacturing Industry. *IOSR Journal of Mechanical and Civil Engineering (IOSRJMCE)*. 2012;1(3):11-14.
 15. Harlisa Mustarudin, Nurani TW. Utilization Rate of Demersal Fish Resources in Pangkep Regency South Sulawesi Indonesia. *Australia and New Zealand Journal of Social Business, Environment and Sustainability*. 2019;5(2).
ISSN 2206-4346
 16. Anderson LG, Juan CS. *Bioeconomics of Fisheries Management Iowa*: Wiley Blackwell Publication. 2010:297.
 17. Auliyah Nurul, Rumagia Faizal, Sinohaji A, Muawanah Umi. Bioeconomic Analysis of Skipjack Tuna Fisheries in North Gorontalo Regency, Indonesia. *IOP Conference Series: Earth and Environmental Science*. 2021;890:012051.
DOI: 10.1088/1755-1315/890/1/012051
 18. Fauzi A, Anna S.. *Modeling of Fisheries and Marine Resources*. Gramedia Pustaka Utama. Jakarta. 2005:343.
 19. Adrianto L. Economic Concepts and Definitions of Coastal and Marine Natural Resources. *PKSPL-IPB Journal*. 2005:1-9.
 20. Bromley DW, Cornea M. *The Management of Common Property Natural Resources*. World Bank Discussion. 1989:57.
 21. Clark CW. *Bioeconomics Modeling and Fisheries Management*. Canada: Vancouver. John Wiley & Sons, Inc. New York. 1985:291.
 22. Csirke J. Small shoaling Pelagic Fish Stock. In Gulland JA (ed.) *Fish Population Dynamics: The Implications for Management*. Chichester. John Wiley & Sons Inc. New York. 1988;422: 271-302.
 23. Gordon HS. The Economic Theory of a Common Property Resource. *The Fishery Journal of Political Economy*. 1954;62:124-142.
 24. Harris JW, Goodwin. *A Survey of Sustainable Development: Social and Economic Dimensions*. The Global Development and Environment. Institute

- Tufts University. Island Press. Washington, Covelo, and London; 2001.
25. Sajeri Harlisa, Mustaruddin Mustaruddin, Nurani Tri. Utilization rate of demersal fish resources in Pangkep Regency South Sulawesi Indonesia. 2019;5:77-82.
 26. Schaefer MB. Some considerations of population dynamics and economics relation to marine fisheries management. Canada: Journal of the Fisheries Research Board. 1957;14:669-681.
 27. Solow R. On the Intertemporal Allocation of Natural Resources. Scandinavian Journal of Economics. 1986;88:141-9.
 28. United Nations Convention on the Law of the Sea. Index and Final Act of the Third United Nations Conference on the Law of the Sea. New York; 1982.

© 2023 *Fidhiani and Rachmawani*; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/97888>