



# **Species of *Monogenean* Parasites Infesting *Cyprinus carpio* and *Planiliza abu* in Lake Balloran Dam-Syria**

**Nada Hamadan <sup>a\*</sup>, Hassan Mohammad Salman <sup>a</sup> and Zouhair Almajid <sup>a</sup>**

<sup>a</sup> *Department of Zoology, College of Sciences, Tishreen University, Lattakia, Syria.*

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/ajfar/2024/v26i11826>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124953>

**Review Article**

**Received: 17/08/2024**

**Accepted: 19/10/2024**

**Published: 28/10/2024**

## **ABSTRACT**

This study was conducted in Lake Balloran Dam and involved 68 fish specimens, including 23 common carp (*Cyprinus carpio*) and 45 Abu mullet (*Liza abu*). Sampling was carried out monthly and randomly from July 2020 to August 2021. The research addressed parasitic infections in these two important species of freshwater fish.

This work identifies five parasitic species of monogeneans, three of them were isolated from common carp and identified as species belonging to the genus *Dactylogyrus*: *D. dogieli*, *D. minutus*, and *D. extensus*. One species was isolated from both the common carp and Abu mullet, and identified as the species *Gyrodactylus derjavini*, and the other species, *G. elegans*, was isolated from only the common carp.

\*Corresponding author: Email: [hamdannada909@gmail.com](mailto:hamdannada909@gmail.com);

**Cite as:** Hamadan, Nada, Hassan Mohammad Salman, and Zouhair Almajid. 2024. "Species of Monogenean Parasites Infesting *Cyprinus Carpio* and *Planiliza Abu* in Lake Balloran Dam-Syria". *Asian Journal of Fisheries and Aquatic Research* 26 (11):23-32. <https://doi.org/10.9734/ajfar/2024/v26i11826>.

Dactylogyrus species were primarily isolated from the gills of the studied fish, whereas Gyrodactylus species were found on the skin and fins.

This study is of fundamental interest for the knowledge of parasitic biodiversity on the Asian continent that are widespread in vast bodies of water, and the species *D. dogieli* and *G. derjavini* were recorded for the first time in Syrian freshwater in this study.

**Keywords:** *Monogenea; Common carp; abu mullet; lake balloran dam.*

## 1. INTRODUCTION

Fish are susceptible to various pathogenic agents such as fungi, viruses, bacteria, and parasites. These parasites can be classified into external parasites, which attach to the outer surface of the fish, and internal parasites, which infest internal organs and muscles, feeding on the fish's internal fluids. This can lead to increased mortality rates in infected fish, directly proportional to the severity of the infestation (Mhaisen, 2019). Monogenean worms are among the most significant and dangerous parasitic worms affecting freshwater fish, due to their direct impact on the tissues of infected fish or their indirect role in secondary bacterial and fungal infections (Al-Saadi and Rashid, 2017). These worms are unique in that they complete their entire life cycle on the fish, requiring no intermediate host, which means they have a direct life cycle. They are also host-specific and show a preference for particular attachment sites. Notable examples include *Dactylogyrus* species, which infest the gills of fish, and *Gyrodactylus* species, which parasitize the skin and fins (Taylor and Francis, 2014).

Few studies have been conducted on wild fish parasites in Syria, with most focusing on parasites in farmed fish. One such study was conducted by Alsamman (Al-Samman, 1989), which investigated the parasites of common carp from several fish farms in both Syria and Hungary. Salman H, and Dayoub, A, (2002), and Salman H, Al-Samman A, Dayoub A, (2003) conducted an ecological and taxonomic study on parasites of common carp in the Al-Sinn River fish farm, isolating and identifying six species of Monogenean worms, including *D. arquatus*, *G. cyprini*, and *G. medius*, which were recorded in Syria for the first time. Salman H, (2004) also studied parasitic ciliates on the same farm. Further research by Alsamman A, Molnar K, Szekely C, (2006) explored Monogenean infestations in both farmed and wild fish in Syria, while Salman H, Al-Samman A, Dayoub A, (2006) identified various species of myxosporean

parasites. Dayoub A, and Salman H, (2015) conducted a study on the use of Monogenean parasites as bioindicators of environmental pollution in Lake 16 Tishreen Dam.

Regionally, several studies have focused on external parasites of mirror carp from the Seyhan River in Turkey (Cengizler et al., 2001). Globally, Martins ML, Onaka EM, Morales FR, Bozzo F, Faro AD, Goncalves A, (2002) conducted research on freshwater fish in various farms across Brazil, where they recorded heavy infestations of external parasites, including Monogenean species.

Based on the previous studies, the objective of this study was to isolate the parasites that are widespread in vast bodies of water in two host fish, the common carp and the Abu mullet, at Lake Balloran Dam which represents a very important freshwater source in the Syrian coastal region.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

Lake Balloran Dam is located in the northwestern part of Syria, near the village of Balloran, to the right of the Latakia-Kassab road, approximately 30 km north of Latakia (35°49'29"N, 35°57'6"E). The dam was built in the valley of the Shamerliya River.

### 2.2 Sample Collection

A total of 68 fish specimens were collected monthly from various sites around the lake between July 20, 2020, and August 15, 2021 (Fig. 1). The fish were captured using gill nets, cast nets, and standard hand fishing rods.

### 2.3 Species Identification and Parasites Isolation

The samples were taken to the Graduate Research Laboratory in the Department of



**Fig.1. Study location at Lake Balloran Dam**

Animal Biology, Faculty of Science, where they were identified to the species level using international taxonomic keys (Beckman, 1962; Berg, 1962; Berg, 1964; Berg, 1965). External examination of the body surface (skin, fins, gills, oral cavity, and nasal pits) of the studied common carp and Abu mullet were studied and the parasites were identified using classification and description studies (Bykhovskaya-Pavlovskaya et al., 1962; Galli et al., 2010; Rahmouni et al., 2023).

### **3. RESULTS AND DISCUSSION**

#### **3.1 Species Identification and Parasites Isolation**

External examination of the body surface of the studied common carp (*Cyprinus carpio*) and Abu mullet (*Liza abu*) revealed the presence of five species of Monogenean parasitic worms. The scientific classification of the isolated species is as follows:

Phylum : Platyhelminthes (Flatworms)  
Class : Monogenea  
Subclass : Polyonchoinea  
Order : Dactylogyridea  
Family : Dactylogyridae  
Genus : Dactylogyrus  
*Dactylogyrus extensus* (Müller & Van Cleave, 1932)  
*Dactylogyrus minutus* (Kulwiec, 1927)  
*Dactylogyrus dogieli* (Gussev, 1953)  
Order : Gyrodactylidea  
Family : Gyrodactylidae  
Genus : Gyrodactylus  
*Gyrodactylus elegans* (Nordmann, 1832)  
*Gyrodactylus derjavini* (Mikailov, 1975)

#### **3.1.1 Parasites of the Genus Dactylogyrus**

Three species belonging to this genus were isolated and identified from the gills of common carp (*Cyprinus carpio*):

##### **3.1.1.1 The species: *D. extensus***

This is a widespread parasite that infests the gills of carp species of various ages. It parasitizes the middle part of the gill lamellae and is relatively large, with a worm length ranging from 580.5 to 1750 microns, averaging 1350.55 microns, and a width ranging from 180 to 490 microns, with an average of 280.95 microns (Fig. A-2).

The hard parts of the attachment disc consist of seven pairs of peripheral hooks, measuring 24-36 microns in length, and one pair of central hooks connected by a linking piece. The copulatory organ consists of a curved projection and a ring-shaped supporting structure, with an average length of 75.15 microns (Fig. B-2).

##### **3.1.1.2 The species: *D. minutus***

This species consists of medium-sized worms, frequently isolated from the base of the gill filaments. The worm length ranges from 315 to 510.15 microns, with an average of 395.085 microns, and a width ranging from 85.90 to 125.50 microns, with an average of 94.80 microns.

The attachment disc is bell-shaped and consists of a pair of large central hooks connected by a dorsal linking piece, and seven pairs of smaller peripheral hooks around the disc's circumference, with an average length of 23.15 microns (Fig. A-3).

The copulatory organ is composed of a long, slender projection, with a gradually tapering base towards the apex, averaging 34.58 microns in length (Fig. B-3).

### 3.1.1.3 The species: *D. dogieli*

This species consists of flatworms that infest the gills of fish. The attachment disc is composed of

seven pairs of small peripheral hooks and one pair of large central hooks connected by a linking piece. There are two pairs of ocular spots at the anterior part of the body (Fig. B-4).

The worm length ranges from 800 to 1050 microns, with an average of 865.75 microns, and a width ranging from 280 to 300 microns, with an average of 283.15 microns (Fig. A-4).




Fig. 2. (A) General appearance of *D. extensus* (x10); (B) Central hooks with the connecting piece (x40)




Fig.3. (A) General appearance of *D. minutus* (x10); (B) Central hooks with connecting piece (x40)






Fig. 4. (A) General appearance of *D. dogieli* (x10); (B) Central hooks with connecting piece (x40)

Table 1. Measurements obtained with a micrometric lens of different body parts in dactylogyrus species parasitic on common carp (*Cyprinus carpio*).

Species + Average Measurements(µm)		<i>D. extensus</i>	$\bar{x}$	<i>D. minutus</i>	$\bar{x}$	<i>D. dogieli</i>	$\bar{x}$
Body	Length	580,5-1750	1350,55	315-510,15	395,85	800-1050	865,75
	Width	180-490	280,95	85,90-125,50	94,80	280-300	283,15
Haptor	Length	80,90-133,65	98,80	58,93-67,22	61,85	47,50-80,95	66,65
	Width	130,15-185,30	95,45	84,50-105,90	93,45	83,16-100,50	97,95
Connecting bar	Length	40,20-51,20	26,10	24,98-26,50	25,81	49-62	56,80
	Width	12,60-15,95	13,75	4,50-5,15	4,95	6-8	7,5
Median hooks	Dorsal length	57,20-74	64,16	34,95-38,15	36,85	4-8	6
	Ventral length	71,80-80,90	73,80	38,90-45	42,80	29-40	35,16
Marginal hooks		31,50-33,60	31,96	18-24	23,15	27,56-34,30	30,25
Copulatory organ		73,50-83,14	75,15	34,15-38	36,58	55,10-72,50	61,35
The infested organ	common carp			the gills		the gills	
	Abu mullet	-		-		-	

### 3.1.2 Parasites of the Genus Gyrodactylus

Two species belonging to this genus were isolated and identified from the skin and fins of common carp (*Cyprinus carpio*) and Abu mullet (*Liza abu*). These species differ in their morphological and anatomical characteristics, as well as in the micrometric (morphological) measurements of various body parts.

#### 3.1.2.1 The species: *G. elegans*

Worms of this species were isolated from the skin of common carp and are relatively large, with a length ranging from 430 to 680 microns, averaging 520.30 microns, and a width ranging from 128.60 to 180.40 microns, with an average of 149.05 microns. The average length of the peripheral hooks is 50.35 microns, and the average total length of the central hooks is 96.93

microns (Fig. A-5). 3.1.2.2 The species: *G. derjavini*

Worms of this species were isolated from the skin, fins, and gills of common carp (*Cyprinus carpio*) and Abu mullet (*Liza abu*). They are relatively large, with a length ranging from 540 to

590 microns, averaging 556.43 microns, and a width ranging from 75.80 to 90.45 microns, with an average of 83.10 microns (Fig. A-6).

The average length of the peripheral hooks is 31.33 microns, and the average total length of the central hooks is 59.83 microns (Fig. B-6).




Fig. 5. (A) General appearance of *G. elegans* (×10); (B) Central hooks with connecting piece (×40)




Fig. 6. (A) General appearance of *G. derjavini* (×10); (B) Peripheral hooks and central hooks with the connecting piece (×40)

**Table 2. Micrometric measurements of different body parts in isolated Gyrodactylus species**

Species + Average Measurements( $\mu\text{m}$ )		G. elegans	$\bar{x}$	G.derjavini	$\bar{x}$
Body	Length	430-680	520,30	540-590	556,43
	Width	128,60-180,40	149,05	75,80-90,45	83,10
Haptor	Length	54,80-63,70	58,30	53,30-64,75	57,90
	Width	47,15-58	51,25	48,20-56,30	49,25
Dorsal connecting bar	Length	21,90-27,50	24,90	19,40-29,25	25,50
	Width	2,80-3,90	3,45	2,10-2,90	2,60
Ventral connecting bar	Length	33,20-43,15	39,15	22,20-28,40	25,75
	Width	7,35-9,90	8,30	7,5-9,30	7,80
Marginal hooks		75,80-135,90	96,93	55,20-63,80	59,83
Median hooks		42-59	50,35	29,50-32,20	31,33
The infested organ	common carp	the skin		the skin, the fins, and the gills	
	Abu mullet	-		the skin, the fins, and the gills	

Parasitic infections result from the interplay of several factors, including those related to the parasite, such as its type and number (severity of infection), and those related to the host, such as its type, sex, age, behavior, and feeding habits, in addition to other environmental factors. Parasitic infestation considers a serious threat for freshwater fishes in tropical and subtropical regions due to severe economic losses either directly or indirectly (Roberts, 1978; Borji et al., 2012). The diversity of parasites in freshwater fish can be attributed to the variation in fish species and the types of parasites present under the same conditions. The appearance of parasites on multiple host fish species indicates that the parasite is widespread, resistant to environmental changes, and adaptable.

Species of the genus *Dactylogyrus* are among the most common parasites of common carp (*Cyprinus carpio*). Our findings are consistent with those of Dayoub, A, (2002) and Zidan M, (2000), who isolated *D. extensus*, *D. minutus*, and *D. dogieli* from the gills of common carp only. The species *D. dogieli* was recorded for the first time in Syrian freshwater, while *D. extensus* and *D. minutus* were recorded by Salman H, and Dayoub, A, (2002) from fish in the Al-San fish farm and Zidan M, (2000) from common carp in Assad Lake. In Iran, more than 70 species of the genus *Dactylogyrus* have been reported from freshwater fishes (Jalali, 1998). The frequency of occurrence of *Dactylogyrus* species which were found as parasites of carp fishes gills in Iran ranged from 2% for the species *D. anchoratus* to

29% to other species like *D. extensus* (Borji et al., 2012).

The morphological and micrometric measurements of the registered *Dactylogyrus* species were consistent with those reported by Oztürk T, & Ozer A, (2014) and Kearns GC, (2014). However, they partially differed from those found by Glaeser HJ, (1969), which may be attributed to differences in surrounding environmental conditions such as temperature, humidity, and dissolved oxygen.

Species of the genus *Gyrodactylus* were primarily isolated from the skin and fins, occasionally from the oral cavity, and rarely from the gills. In general, *Gyrodactylus* species are the most common skin parasites infecting freshwater fishes all over the world and causing serious diseases to the common carp (Yamaguti, 1963; Abdullah and Mama, 2013). This finding is in agreement with Woo PTK, (2006) and Al-Zubaidy A, (2007). *G. derjavini* was recorded for the first time in Syria in our study, while *G. elegans* was recorded by Salman H, Al-Samman A. Dayoub A, (2003) from common carp in the Al-San fish farm and Safar G, (2019) from common carp in the Salah al-Din Dam (Safarqiya). In Iraqi Kurdistan, several species of *Gyrodactylus* were recorded in the gills, fins, and skin of the common carp (Abdullah and Mama, 2013).

The morphological and micrometric characteristics of the registered *Gyrodactylus* species were consistent with measurements reported by Gussev AV, (1985) and Malmberg G,

(1970). However, they partially differed from those reported by Al-Zubaidy A, (2007), likely due to differences in the fish species from which the parasites were isolated; in our study, parasites were isolated from Abu mullet and common carp, whereas Al-Zubaidy A, (2007) isolated them from *Aspius vorax*. In other inventory in Iraqi Kurdistan, 10 species of the genus *Dactylogyrus* and 8 species of the genus *Gyrodactylus* were recorded parasiting on common carp, *Dactylogyrus* species were found in the gills and *Gyrodactylus* species were found in the gills and skin (Mama and Abdullah, 2012).

#### 4. CONCLUSIONS

The results of this study indicate that species of the genus *Dactylogyrus* primarily parasitize the gills of fish, while species of the genus *Gyrodactylus* are found on the skin and fins. Notably, *G. derjavini* and *D. dogieli* were recorded for the first time in the freshwater of the Syrian coast in this study.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Authors hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

Abdullah, S. M., & Mama, K. S. (2013). Parasitic infections with *Gyrodactylus* (Monogenea) on common carp (*Cyprinus carpio*) from Ainkawa fish hatchery in Erbil city, Kurdistan Region, Iraq. In *Proceedings of the 4th ICOWOBAS-RAFSS*, Johor Bahru, Malaysia, 3-5.

Al-Saadi, A. A., & Rashid, R. (2017). First record of three monogenean species of the genus *Gyrodactylus* from the gills of some Tigris River fish. *Ibn Al-Haitham Journal for Pure and Applied Sciences*, 30(1), 306-3014.

Al-Samman, A. (1989). Incidence of Monogenean species on the gill of common carp (*Cyprinus carpio*) collected

from Hungarian and Syrian fish farms. University of Agricultural Sciences. 45 – 50

Al-Samman, A., Molnar, K., & Szekely, C. (2006). Infection of cultured freshwater fish with monogeneans in Syria. *Bulletin of the European Association of Fish Pathology*, 26(4), 170-173.

Al-Zubaidy, A. (2007). First record of three monogenic parasite species from Iraqi freshwater fishes. *JKAU: Mar. Sci.*, 18, 83-94.  
Available: <https://doi.org/10.30526/30.1.1079>

Beckman, W. C. (1962). The freshwater fishes of Syria. FAO Fisheries Biology Technical, 8, 126-133.

Berg, L. S. (1962). *Freshwater fishes of the U.S.S.R. and adjacent countries* (Vol. I). Academy of Sciences of the U.S.S.R. Zoological Institute.

Berg, L. S. (1964). *Freshwater fishes of the U.S.S.R. and adjacent countries* (Vol. II). Academy of Sciences of the U.S.S.R. Zoological Institute.

Berg, L. S. (1965). *Freshwater fishes of the U.S.S.R. and adjacent countries* (Vol. III). Academy of Sciences of the U.S.S.R. Zoological Institute.

Borji, H., Naghibi, A., Nasiri, M. R., & Ahmadi, A. (2012). Identification of *Dactylogyrus* spp. and other parasites of common carp in northeast of Iran. *Journal of Parasitic Diseases*, 36, 234-238.

Bykhovskaya-Pavlovskaya, I. E., Gusev, A. V., Dubinina, M. N., Izyumova, N. A., Smirnova, T. S., Sokolovskaya, I. L., Shtein, G. A., Shul'man, S. S., & Epshtein, V. M. (1962). *Key to parasites of freshwater fish of the USSR*. Akad. Nauk, SSSR Moscow, 727

Cengizler, I., Aytac, N., Azizoglu, S. A., & Ozak, A. A. (2001). Ecto-endo parasite investigation on mirror carp (*Cyprinus carpio* L., 1758) captured from the river Seyhan, Turkey. *E.U. Journal of Fisheries & Aquatic Sciences*, 18(1), 87-90.

Dayoub, A. (2002). Environmental and taxonomic study of some freshwater fish parasites in the Syrian Coastal Region (Master's thesis, Department of Zoology, Faculty of Science, University of Tishreen, Syria). [In Arabic].



- Dayoub, A., & Salman, H. (2015). Study of using Monogenea parasites on free-living fishes in the Lake of 16 Tishreen Dam as bioindicators of environment pollution. *International Journal of Biomedical Engineering and Clinical Science*, 1(1), 15-22.  
DOI:10.4194/1303-2712-v14\_2\_07
- Galli, P., Pugachev, O., & Kritsky, D. (2010). *Guide to Monogenoidea of freshwater fish of Palaearctic and Amur regions*.
- Glaeser, H. J. (1969). Die Gyrodactylus-Arten des Karpens in der Deutschen Demokratischen Republik. *Deutsche Fischerei-Zeitschrift*, 16, 372-380.
- Gussev, A. V. (1985). *Key of freshwater fish parasites*. Institute of Zoology, Academy of Science, Leningrad: USSR, 425.
- Jalali, B. (1998). *Parasites and parasitic diseases of freshwater fishes of Iran*. Fisheries Co of Iran.
- Kearn, G. C. (2014). Some aspects of the biology of monogenean (Platyhelminth) parasites of marine and freshwater fishes. *Oceanography*, 2, 117.
- Malmberg, G. (1970). The excretory system and the marginal hook as a basis for the systematics of *Gyrodactylus* (Trematoda, Monogenea). *Arkiv für Zoologie*, 23(1), 1-235.
- Mama, K. S., & Abdullah, S. M. (2012). A comparative study on the parasitic fauna of the common carp (*Cyprinus carpio*) from Ainkawa fish hatchery (Erbil) and Lesser Zab river in Kurdistan region, Iraq. *Mesopotamian Journal of Agriculture*, 42(2), 19-26.
- Martins, M. L., Onaka, E. M., Morales, F. R., Bozzo, F., Faro, A. D., & Goncalves, A. (2002). Recent studies on parasitic infections of freshwater cultivated fish in the state of São Paulo, Brazil. *Maringá*, 24(4), 981-985.
- Mhaisen, F. T. (2019). Checklists of parasites of fishes of Thi-Qar province, Iraq. *Biological Applied Environmental Research*, 3(2), 152-167.
- Öztürk, T., & Özer, A. (2014). Monogenean fish parasites, their host preferences and seasonal distributions in the Lower Kızılırmak Delta (Turkey). *Turkish Journal of Fisheries and Aquatic Sciences*, 14, 367-378.
- Rahmouni, C., Seifertová, M., Benovics, M., & Šimková, A. (2023). Diversity and phylogeny of *Gyrodactylus* spp. (Monogenea: Gyrodactylidae) across the Strait of Gibraltar: Parasite speciation and historical biogeography of West Mediterranean cyprinid hosts. *Diversity*, 15(11), 1152.
- Roberts, R. J. (1978). *Fish pathology*. Bailliere Tindall.
- Safar, G. (2019). Study of the species composition of monogenean parasites on fish in the Salah al-Din Dam (Safarqiya) (Master's thesis, Department of Animal Life Sciences, Faculty of Science, University of Tishreen, Syria). [In Arabic].
- Salman, H. (2004). Contribution to the study of some ciliated parasites (Protozoa) in common carp (*Cyprinus carpio* L.) in the Al-San Fish Farm. *Tishreen University Journal for Scientific Research (Basic Sciences Series)*, 26(3), 111-121.
- Salman, H., & Dayoub, A. (2002). Dynamics of infection by *Dactylogyrus* worms parasitizing carp in the Al-San Fish farm. *Tishreen University Journal for Scientific Studies and Research*, 24(12), 159-172.
- Salman, H., Al-Samman, A., & Dayoub, A. (2003). *Gyrodactylus* spp. worms parasitizing carp in the Al-San Fish Farm. *Tishreen University Journal for Scientific Studies and Research*, 25(13), 145-157.
- Salman, H., Al-Samman, A., & Dayoub, A. (2006). First record of radiolarian sporozoans (gelatinous animals) in ciliated worms in freshwater of the Syrian Coastal Region. *Tishreen University Journal for Scientific Studies and Research*, 28(2), 175-184.
- Taylor & Francis. (2014). *Parasitic worms of fish*. Ltd. 443
- Woo, P. T. K. (2006). *Fish diseases and disorders* (Vol. 1): Protozoan and Metazoan infections (2nd ed.). University of Guelph Canada. 802,
- Yamaguti, S. (1963). *Systema helminthum* (Vol. IV). Monogenea and Aspidocotylea.

Zidan, M. (2000). Study of the distribution of parasitic worms in common carp (*Cyprinus carpio* L.) in Assad Lake (Master's thesis,

Department of Zoology, Faculty of Science, University of Aleppo, Syria), 175.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

---

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/124953>