



## **Incidence of Esophageal Varices in Patients with Upper Gastrointestinal Bleeding**

**Khalidurehman Basharat<sup>a#</sup> and Wali Gul Mokhles<sup>a\*</sup>**

<sup>a</sup> Faculty of Medicine, Shaikh Zayed University, Khost, Afghanistan.

### **Authors' contributions**

*This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/JPRI/2022/v34i19A35804

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/82999>

**Received 30 December 2021**

**Accepted 01 March 2022**

**Published 07 March 2022**

**Original Research Article**

## **ABSTRACT**

**Objective:** To determine the frequency of esophageal varices as a cause of upper gastrointestinal bleeding.

**Methodology:** This descriptive cross sectional study was conducted in the emergency room and the department of medicine Khost Provincial Hospital Khost -Afghanistan. Duration of data collection was six months, starting from January 5, 2021 to June 4, 2021. The study included one hundred and six patients suffered from upper GI bleeding. After initial history and clinical examination, upper GI endoscopy was performed to evaluate the cause of bleeding and all the relevant data was entered in the specific proforma designed by researchers.

**Results:** One hundred and six patients, 60 (56.6%) male and 46 (43.4%) females, of upper GI hemorrhage were included in the study. Mean age was 45.78 years (SD  $\pm$ 14.86) with age range of 21- 85years. Twenty patients (18.86%) had esophageal varices while 28 (26.41%) patients had duodenal ulcer, 31 (29.2%) had gastric ulcer, 14 (13.2%) had Gastric carcinoma, 9 (8.49) patients had gastro-duodenal erosions, three patients (2.83%) had Mallory Weiss tear and one patient (0.94) had esophageal growth.

**Conclusion:** In this area (Khost – Afghanistan) where this study was conducted, esophageal varices are the second most common cause of upper GI bleeding.

<sup>#</sup> Senior Lecturer;

<sup>\*</sup> Assistant Professor;

\*Corresponding author: E-mail: walimokhles@yahoo.com;

**Keywords:** Duodenal ulcer; endoscopy; esophageal varices; gastrointestinal hemorrhage; gastric ulcer hematemesis; incidence; liver cirrhosis; melena.

## 1. INTRODUCTION

Upper gastrointestinal (GI) bleeding is a common reason for hospital admission in older adults and carries a high morbidity and mortality if not properly managed. Both hematemesis and melena or either alone, are famous manifestations of Bleeding from gastrointestinal tract [1]. Upper GI hemorrhage may rarely present in the form of hematochezia [2].

The definition of hematemesis is vomiting of gross blood [1]. In United States, the annual incidence of upper GI hemorrhage is approximately 48-160/100,000 with a mortality of 5% [3]. Increasing age, co-morbidities and hemodynamic compromise (tachycardia or hypotension) are three independent clinical predictors of death in patients hospitalized for upper GI hemorrhage [4].

A number of sources that upper GI hemorrhage may originate from, are:

1. Esophageal varices & portal hypertensive gastropathy.
2. Peptic ulcer disease.
3. Gastric neoplasms.
4. Gastro-duodenal erosions
5. Erosive esophagitis.
6. Mallory Weiss tear.
7. Others as angiodysplasia, Dieulafoy's lesion, aortoenteric fistula, hemobilia, hereditary hemorrhagic telangiectasia, uremia and coagulation disorders [5].

Between different geographical areas of the world, the multifactorial etiology of upper GI bleeding varies widely. Peptic ulcer and esophageal varices are the two most common causes of the GI bleeding and of the two the esophageal varices are the most common cause of upper GI bleeding in India [6]. Esophageal varices are dilated submucosal veins that build up in patients with causal portal hypertension. Cirrhosis of liver is the most common cause of portal hypertension. More recent data propose that the amount of cases as a consequence of peptic ulcer disease has decreased [7].

Variceal bleeding is the cause of bleeding in cirrhotic patients in 50-60%. Rebleeding in upper gastrointestinal bleeding occurs in 7-16%, despite endoscopic therapy. Re-bleeding is

especially high in variceal bleeding and peptic ulcer bleeding. Mortality ranges between 3 and 14% and didn't change in the past 10 years [8]. Mortality risk is about 24.1 % after each episode of variceal bleeding. Until gastro-esophageal varices are obliterated, the risk of re-bleeding is about 29.4% high [9]. After variceal hemorrhage, one year survival can be poor. The severity of the liver disease, the size of the varices and the presence of red markings on the varices are considered the three independent predictors of bleeding [10].

The study which was conducted, to find out the frequency of esophageal varices in upper GI bleeding; a common medical emergency in Khost province of Afghanistan. The potentially fatal complications of chronic hepatitis and cirrhosis like esophageal varices are reduced by avoidance and on time treatment of the mentioned diseases.

## 2. METHODOLOGY

This is a descriptive cross sectional study. It was conducted in the emergency room and the department of medicine in Khost Provincial Hospital Khost -Afghanistan. The data was collected in six months, starting from January 5, 2021 to June 4, 2021. One hundred six patients of upper GI bleeding were included in the study. Nonprobability purposive sampling technique was used. All the patients who were adult coming to the hospital with upper gastrointestinal bleeding in the form of hematemesis or melena, were included. Exclusion criteria were: 1. when in the upper or lower airway, there was a source of bleeding. 2. Non-cooperative patient as those with the advanced hepatic encephalopathy. 3. Because of some medical problems such as severe cardiac or respiratory illness or hemodynamically unstable patient was not fit for endoscopy. 4. Patients who rejected hospital admission for further investigations and treatment.

During the data collection period of six months, totally one hundred six (106) patients of upper GI bleeding were included in the study. The patients of upper GI bleeding on the basis of hematemesis and melena, were diagnosed clinically. These patients were admitted through the emergency room and the medical outdoors of Khost Provincial Hospital Khost – Afghanistan.

Regarding admission and further treatment, informed consent was taken. Taking history, which included the duration and amount of hematemesis and color of the vomitus, made the provisional diagnosis. About precipitating factors such as alcohol taking, NSAIDs use or the use of anticoagulants was asked. Inclusive general physical examination was performed. Digestive, respiratory, cardiovascular and central nervous system were examined along with examination for epigastric tenderness and signs of chronic liver disease. Upper GI endoscopy performed which served both diagnostic and therapeutic purposes. Endoscopic evaluation of patients with upper GI bleeding was done by the endoscopy unit of Lemar diagnostic and therapeutic Center Khost city – Afghanistan. Procedure was done using 4% Xylocaine throat spray. No sedating premedication was used. For growth and gastric ulcer with doubt of malignancy, biopsies were taken from growth and the ulcer periphery and were sent for histopathology. For analyzing of data, Statistical Package for Social Science (SPSS) version 20 was used. Descriptive statistics for numerical data i.e. age was calculated as mean  $\pm$  SD, while categorical data i.e. gender, esophageal varices and other causes of upper GI bleeding were calculated as frequencies and percentages. Test of significance was not used and level of significance (p- value) was not estimated, because this was a descriptive study and no need for p- value estimation.

### 3. RESULTS

The study included a total of one hundred six,60 (56.6%) male and 46 (43.4%) female patients. Mean age was 45.78 years (SD  $\pm$ 14.86) with age range of 21- 85 years. Fifty nine (55.66%) patients presented with both hematemesis and melena while 25 (23.59%) patients had only melena and 18 (16.98%) patients had hematemesis alone. Twenty (18.86%) patients 12 male and 8 female had variceal bleeding. The

provisional etiological diagnosis of cirrhosis of liver was confirmed in all patients with variceal bleeding. Jaundice was present in 12 (60%) patients. All patients had one or other signs of cirrhosis such as abdominal distention/ascites, splenomegaly, spider nevi, gynecomastia, decreased area of liver dullness and loss of normal distribution of hair. Viral markers serology showed that 9 (45%) patients were positive for anti HCV antibodies and 8 (40%) patients were positive for hepatitis B surface antigen. In the remaining 03 (15%) patients, viral markers were negative. One of them had Wilson’s disease and two were non-alcoholics steatohepatitis. Two patients had portal hypertensive gastropathy along with esophageal varices. Tachycardia, hypotension and pallor were detected in 9(45%) patients while the remaining were hemodynamically stable.

Gastric ulcer was diagnosed in 31 patients (29.2%), 17 males and 14 females. Epigastric pain history was present in 16 patients. Five patients suffered from mild epigastric tenderness. Eight patients (5 males & 3 female) were using aspirin for osteoarthritis. In six patients, upper GI bleeding caused hypotension and tachycardia.

Twenty eight (26.41%) patients were suffering from duodenal ulcer (15 males; 13 females). The history of duodenal ulcer with hunger pains and relief of symptoms with food and antacids was present in 19 patients, while the other patients didn’t give such history. Twenty patients were positive for stool H.pylori antigen. None of the patients was alcoholic. Hypotension, pallor and tachycardia were present in six patients while the remaining were hemodynamically stable.

Nine (6 male 3 female) had Acute gastroduodenal erosions. Seven (7) patients had the history of NSAIDs ingestion for various reasons. Onset of hematemesis was sudden with mild epigastric pain and tenderness.

**Table 1. Shows the serology for viral markers**

Type of virus	Numbers	Results	%
HBs Ag	8	Positive	40
	12	Negative	60
Anti HCV	9	Positive	45
	11	Negative	55

**Table 2. Endoscopic diagnosis of patients suffered from upper gastrointestinal hemorrhage**

Serial No.	Causes of upper GI hemorrhage	Male	Female	Total	Percentage
1	Esophageal varices	12	8	20	18.86%
2	Gastric ulcers	17	14	31	29.2%
3	Duodenal ulcers	15	13	28	26.41%
5	Mallory Weiss tear	1	2	3	2.83%
6	Gastric carcinoma	8	6	14	13.2%
7	Esophageal growth	1	0	1	0.94
8	Gastro duodenal erosions	6	3	9	8.49
Total		60	46	106	100

Peptic ulcer and esophageal varices are accounted the two most common causes of the gastrointestinal bleeding. Gastric carcinoma was the third common cause of upper GI bleeding in this study, but it is said that variceal bleeding has been found a relatively unusual cause of upper GI bleeding in the Western countries 5.1% [11]. In the United States, among the total cases of upper GI bleeding, the percentage of variceal bleeding varies from 4% to 20% [12] and bleeding due to peptic ulcers account higher than 50% [13]. But in the region, for example in Pakistan , the Variceal bleeding revealed the most common cause of upper GI bleeding in several studies [14,15]. A Gastric growth (Carcinoma of the stomach) was responsible for upper gastrointestinal bleeding in 14 patients (13.2%). These patients were more than 40 years old and had a history of hematemesis, melena, loss of weight, epigastric pain and anemia. Clinical examination showed marked weight loss and mild epigastric tenderness. Endoscopy revealed gastric ulcers with suspicion of malignancy and growths which were proved by histopathology. One patient had esophageal growth suffering from dysphagia and hematemesis.

Mallory Weiss tear was responsible for upper gastrointestinal bleeding in three patient (2.83%).

#### 4. DISCUSSION

Acute upper gastrointestinal bleeding (UGIB) continues to be a common cause of hospital admission and morbidity and mortality [16] The most common etiology of upper GI bleeding is peptic ulcer disease, followed by erosive gastroesophagitis, gastroesophageal varices, Mallory-Weiss tears, Dieulafoy's lesions, gastric antral vascular ectasia (GAVE), portal hypertensive gastropathy (PHG), aortoenteric fistula, malignancy, and other rare conditions [17]. The primary diagnostic test for evaluation of

upper GI bleeding is endoscopy. Endoscopy for upper GI bleed has a sensitivity of 92%–98% and specificity of 30%–100% [18].

Peptic ulcer and esophageal varices are accounted the two most common causes of the gastrointestinal bleeding. Gastric carcinoma was the third common cause of upper GI bleeding in this study, but it is said that variceal bleeding has been found a relatively unusual cause of upper GI bleeding in the Western countries 5.1% [11]. In the United States, among the total cases of upper GI bleeding, the percentage of variceal bleeding varies from 4% to 20% [12] and bleeding due to peptic ulcers account higher than 50% [13]. But in the region, for example in Pakistan , the Variceal bleeding revealed the most common cause of upper GI bleeding in several studies [14,15]. A study conducted in Nepal about the causes of upper gastrointestinal bleeding, revealed that esophageal varices were (47.5%) and peptic ulcer disease (33.3%) [19]. A study which was completed on 114 patients in 2019 in Himalaya medical institute in India, outlined some of the most important causes of bleeding from upper gastrointestinal tract with gastric and esophageal varices being the most significant (56.14%) followed by peptic ulcer (14.91%) [6]. The reasons that in our study the variceal bleeding was the second most common cause of gastrointestinal bleeding are :

1. In Pakistan Meta-analyses estimated the pooled mean HCV prevalence at 6.2% among the general population, 34.5% in high-risk clinical populations, 12.8% in populations at intermediate risk, 16.9% among special clinical populations, 55.9% among populations with liver-related conditions and 53.6% among people who inject drugs [20]. In Afghanistan.

Meta-analyses estimated HCV prevalence at 0.7% among the general population, 32.6% among PWID and 2.3% among populations at intermediate risk [21].

2. The prevalence of *H.pylori* is another reason that percentage of peptic ulcer is higher than esophageal varices as a cause of GI bleeding. A study conducted in Pakistan, total of 201 (45%) cases were found positive for *H. pylori* out of 450 dyspeptic individuals [22] while in Afghanistan, 152 patients with dyspepsia were included in a study; 59 were men (40.0%), and 93 were women (60.0%). The overall seroprevalence of *H. pylori* was 75.6% [23].

But the result of our study is consistent with results of Iranian and Saudi Arab studies. A study which was completed in Shiraz medical university Iran in 2009 on 572 patients showed that 44% of bleeding from upper gastrointestinal tract was caused by Peptic ulcer [24]. As well as , in Saudi Arabia found in a study that was conducted in Riyadh (King Saud Medical City) that Nonvariceal upper GI bleeding source was 80.5% and variceal source was found 13.1% while no endoscopic findings were found in 6.4% of cases [25].

## 5. CONCLUSION

It is worth mentioning to realize that gastrointestinal bleeding is a fatal and challenging problem and caused mostly by peptic ulcer and esophageal varices in Afghanistan. Esophageal variceal bleeding is a shocking complication of portal hypertension and is an important cause of disability and death in patients with upper GI bleeding. Once variceal bleeding has occurred, the prognoses are poor, so prevention of chronic hepatitis and cirrhosis is extremely crucial.

## CONSENT

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

## ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Costable NJ, Greenwald DA. Upper gastrointestinal bleeding. Clin Geriatr Med. 2021 Feb;37(1):155-172. DOI: 10.1016/j.cger.2020.09.001.PMID:33213769.
2. Ray-Offor E, Elenwo S. Endoscopic evaluation of upper and lower gastrointestinal bleeding. Nigerian Journal of Surgery. 2015;21(2):106. DOI: 10.4103/1117-6806.162575
3. Tielleman T, Bujanda D, Cryer B. Epidemiology and risk factors for upper gastrointestinal bleeding. Gastrointestinal Endoscopy Clinics of North America. 2015;25(3):415-428. DOI: 10.1016/j.giec.2015.02.010
4. Khamaysi I, Gralnek IM. Acute upper gastrointestinal bleeding (UGIB) – Initial evaluation and management. Best Practice & Research Clinical Gastroenterology. 2013;27(5):633-638. DOI: 10.1016/j.bpg.2013.09.002
5. Feinstein LB, Holman RC, Christensen KLY, Steiner CA, Swerdlow DL. Trends in hospitalizations for peptic ulcer disease, United States, 1998-2005. Emerging Infectious Diseases. 2010;16(9):1410-1418. DOI: 10.3201/eid1609.091126
6. Deep Anand, Rohit Gupta, Minakshi Dhar, Vivek Ahuja Department of medicine, himalayan institute of medical sciences, 1department of medicine, unit of gastroenterology, Himalayan Institute of Medical Sciences, Swami Ram Nagar, Jolly Grants, Doiwala, Dehradun, Uttarakhand, India.
7. Hilzenrat N, Sherker AH. Esophageal varices: Pathophysiology, approach, and clinical dilemmas. International Journal of Hepatology. 2012;1-2. DOI: 10.1155/2012/795063
8. VanLeerdam ME. Epidemiology of acute upper gastrointestinal bleeding. Best

- Practice & Research Clinical Gastroenterology. 2008;22(2):209–224.  
DOI: 10.1016/j.bpg.2007.10.011
9. Olmo JA de, Peña A, Serra MA, Wassel AH, Benages A, Rodrigo JM. Predictors of morbidity and mortality after the first episode of upper gastrointestinal bleeding in liver cirrhosis. *Journal of Hepatology*. 2000;32(1):19–24.  
DOI: 10.1016/s0168-8278(01)68827-5
  10. Ferguson JW, Tripathi D, Hayes PC. Endoscopic diagnosis, grading and predictors of bleeding in esophageal and gastric varices. *Techniques in Gastrointestinal Endoscopy*. 2005;7(1):2–7.  
DOI: 10.1016/j.tgie.2004.10.002
  11. Hreinsson JP, Kalaitzakis E, Gudmundsson S, Björnsson ES. Upper gastrointestinal bleeding: incidence, etiology and outcomes in a population-based setting. *Scandinavian Journal of Gastroenterology*. 2013;48(4):439–447.  
DOI: 10.3109/00365521.2012.763174
  12. Tiellemann T, Bujanda D, Cryer B. Epidemiology and risk factors for upper gastrointestinal bleeding. *Gastrointestinal Endoscopy Clinics of North America*. 2015;25(3):415–428.  
DOI: 10.1016/j.giec.2015.02.010
  13. Boonpongmanee S, Fleischer DE, Pezzullo JC, Collier K, Mayoral W, Al-Kawas F, Benjamin SB. The frequency of peptic ulcer as a cause of upper-GI bleeding is exaggerated. *Gastrointestinal Endoscopy*. 2004;59(7):788–794.  
DOI: 10.1016/s0016-5107(04)00181-6
  14. Shah Syed Muhammad Ali, et al. Etiology of upper gastrointestinal bleed at Aziz Bhatti Shaheed Teaching Hospital Gujrat. *Annals of PIMS ISSN 1815*. 2016;2287.
  15. Sher Farrukh, et al. Frequency of different causes of upper gastrointestinal bleeding using endoscopic procedure at a tertiary care hospital. *PAFMJ* 64.3. 2014;410-3.
  16. Ghoddoosi M, Jazi K, Hajrezaei Z, Habibi MA, Ahmadpour S, Modares MP. Upper gastrointestinal bleeding due to concomitant esophageal varices and herpes simplex virus esophagitis in a 70-year-old patient. *Gastroenterology Nursing*. 2021 Sep 1;44(5):357-60.
  17. Costable NJ, Greenwald DA. Upper gastrointestinal bleeding. In: Pitchumoni C.S., Dharmarajan T. (eds) *Geriatric Gastroenterology*. Springer, Cham; 2020.  
Available: [https://doi.org/10.1007/978-3-319-90761-1\\_47-1](https://doi.org/10.1007/978-3-319-90761-1_47-1)
  18. Mahajan P, Chandail VS. Etiological and endoscopic profile of middle aged and elderly patients with upper gastrointestinal bleeding in a Tertiary Care Hospital in North India: A retrospective analysis. *J Midlife Health*. 2017;8(3):137-141.  
DOI: 10.4103/jmh.JMH\_86\_17
  19. Dewan KR, Patowary BS, Bhattarai S. A study of clinical and endoscopic profile of acute upper gastrointestinal bleeding. *Kathmandu Univ Med J*. 2014;45(1): 21-25.
  20. Abu-Raddad Laith J. The epidemiology of hepatitis C virus in Pakistan: Systematic review and meta-analysesR. *Soc. open sci*. 2018;5180257180257.  
Available: <http://doi.org/10.1098/rsos.180257>
  21. Chemaitelly H, Mahmud S, Rahmani AM, Abu-Raddad LJ. The epidemiology of hepatitis C virus in Afghanistan: systematic review and meta-analysis. *International Journal of Infectious Diseases*. 2015;40:54–63.  
DOI: 10.1016/j.ijid.2015.09.011
  22. Khan A, Farooqui A, Raza Y, Rasheed F, Manzoor H, Akhtar SS, Quraishy MS, Rubino S, Kazmi SU, Paglietti B. Prevalence, diversity and disease association *Helicobacter pylori* in dyspeptic patients from Pakistan. *J Infect Dev Ctries*. 2013;7:220–228.  
DOI: 10.3855/jidc.2942
  23. Hamrah MH, Hamrah MS, Hassan Hamrah M, et al. Prevalence of *Helicobacter pylori* infection in dyspeptic patients in andkhoy afghanistan. *Asian Pac J Cancer Prev*. 2017;18(11):3123-3127.  
Published 2017 Nov 26.  
DOI: 10.22034/APJCP.2017.18.11.3123
  24. Pirastehfar M, Azari A, Saberifiroozi M, Kaviani M. Etiology and outcome of patients with upper gastrointestinal bleeding: A study from South of Iran. *Saudi Journal of Gastroenterology*. 2010;16(4):253.  
DOI: 10.4103/1319-3767.70608

25. Alruzug IM, Aldarsouny TA, Semaan T, Aldaher MK, AlMustafa A, Azzam N, Aljebreen A, Almadi MA. Time trends of causes of upper gastrointestinal bleeding and endoscopic findings. Saudi J of Gastroenterol: 2021;27(1):28.-34.  
DOI:10.4103/sjg.SJG\_378\_20.PMID:3307820;PMCID:PMC8083243

---

© 2022 Basharat and Mokhles; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

*<https://www.sdiarticle5.com/review-history/82999>*